

Quantum measurement corrections to chemically induced dynamic nuclear polarization

I. K. Kominis

Department of Physics, University of Crete, Heraklion 71103, Greece

Chemically induced dynamic nuclear polarization has emerged as a universal signature of spin order in photosynthetic reaction centers. Such polarization, significantly enhanced above thermal equilibrium, is known to result from the nuclear spin sorting inherent in the radical pair mechanism underlying long-lived charge-separated states in photosynthetic reaction centers. We will here show that the recently understood fundamental quantum dynamics of radical-ion-pair reactions open up a new and completely unexpected venue towards obtaining CIDNP signals. The fundamental decoherence mechanism inherent in the recombination process of radical pairs is shown to produce nuclear spin polarizations on the order of 10^4 times or more higher than thermal equilibrium values at low fields relevant to natural photosynthesis in earth's magnetic field. This opens up the possibility of a fundamentally new exploration of the biological significance of high nuclear polarizations in photosynthesis.

The quantum dynamics of photosynthesis have, quite naturally, attracted a lot of attention recently. Understanding if and how Nature exploits quantum (de)coherence would have tremendous scientific and technological impact. For what follows, photosynthesis can be simply visualized as a two-step process: (i) light harvesting and (ii) photochemical processing. Light harvesting molecules absorb the incident photon and guide the subsequent excitation to the photosynthetic reaction center (RC). It is there where the photochemistry takes place, transforming the initial electronic excitation in a transmembrane proton pump that further drives the life-sustaining chemistry. Quantum coherence effects in light harvesting complexes (time scale of 1-500 fs) have been addressed theoretically and experimentally [1-9]. We will here focus on RC quantum dynamics (time scale of 1ps - 10 μ s) schematically depicted in Fig. 1A. Photosynthetic reaction centers exhibit a cascade of electron transfer reactions, starting from a photo excited donor-acceptor dyad PI (chlorophyll-type molecules) and shelving the electron further apart to quinones (Q_A , Q_B), producing a long-lived charge-separated state driving the transmembrane proton pump. In each of those steps there is a radical-ion pair formed, further perplexing the dynamics, as radical-ion pairs (RPs) exhibit intricate magnetic field effects [10]. Radical-ion pairs [11, 12] are biomolecular ions with two unpaired electrons (denoted by the two dots in Fig. 1) and any number of magnetic nuclei. The magnetic nuclei of the donor and acceptor molecules couple to the two electrons via the hyperfine interaction, leading to singlet-triplet (S-T) mixing, i.e. a coherent oscillation of the total electron spin state, also affected by the electrons' Zeeman interaction with the external magnetic field. A spin-dependent charge recombination terminates the coherent S-T mixing and leads to the reaction products, either the original neutral dyad PI (singlet product) or a triplet intermediate (^{3} PI). Singlet (triplet) radical pairs recombine at the singlet (triplet) recombination rate, denoted by k_S (k_T). Triplet products may close the reaction cycle by intersystem-crossing into the

singlet diamagnetic ground state at a rate k_{isc} .




FIG. 1: (A) Cascade of electron transfer reactions taking place in photosynthetic reaction centers. (B) Each charge-separated state in (A) is a radical-ion pair which undergoes additional quantum dynamics due to internal magnetic interactions, embodied by the Hamiltonian $\mathcal{H}$, of the two unpaired electrons with each-other, the magnetic nuclei and the external magnetic field. These magnetic interactions can give rise to singlet-triplet conversion during the finite lifetime of the charge-separated state, further perplexing the dynamics of the whole reaction. Figure adapted from the review of Boxer, Chidsey and Roelofs [10].

It is apparent from Fig. 1 that when the lifetime of the charge-separated states (which increases with separation) is such that magnetic-field effects (with typical timescale of e.g. S-T mixing on the order of 0.1-1 μ s) can come into play, radical-pair dynamics get convolved into the reaction center dynamics. One aspect of this

interrelation pervading a large number of photosynthetic reaction centers is chemically induced dynamic nuclear polarization [13-20], the enhancement (by several orders of magnitude) of the ground state (PI) nuclear spin polarization resulting from the RP spin dynamics. CIDNP has emerged as a rather universal signature of spin order in photosynthetic reaction centers [21] and its possible operational significance for photosynthesis is an open fundamental question. CIDNP is based on nuclear spin sorting, to be shortly explained, taking place in radical-pair reactions. Three fundamental mechanisms have so far been known to produce CIDNP signals, the three-spin-mixing (TSM) [22], the differential decay (DD) and the differential relaxation (DR) mechanism [23]. Note that the first is a coherent mechanism resulting from particular magnetic interactions within the RP, while the other two are incoherent mechanisms, of which the second (DR) will not be relevant for the following discussion. We will here show that there is yet another, completely unanticipated mechanism at work *at the fundamental quantum dynamical level of RP reactions*. It is due to the inherent quantum measurement of the RP's spin state continuously going on in the RP as part of its very charge recombination process. The decoherence caused by this "internal" quantum measurement is responsible for a new kind of nuclear spin sorting, particularly efficient at low fields, and completely unrelated to other low-field CIDNP studies [24]. This new mechanism is ubiquitous, as it is largely independent of the particular magnetic interactions in the radical pairs, and thus opens a new venue towards exploring the biological significance of CIDNP-induced spin order in natural (earth's field) photosynthesis. In the following we will (i) briefly explain the basics of CIDNP signal generation for the general reader and (ii) recapitulate the recent progress in understanding the fundamental quantum dynamics of radical-ion-pair reactions. Combining (i) and (ii), we will then explain the quantum measurement corrections to CIDNP.

Consider a radical-ion pair with just one nucleus coupled to the electron of e.g. the donor. At time $t = 0$ the radical-ion pairs start out in the electronic singlet state, while the nuclear spins are thermally polarized, which for protons at earth's field corresponds to a polarization of about 10^{-10} , practically taken to equal zero. Thus, in the ensemble of radical-ion pairs half will have the nuclear spin in the $|+\rangle$ state, while in the other half the nuclear spin will initially be in the $|-\rangle$ state. The local magnetic field seen by the donor electron will be the external field plus the nuclear magnetic field, while the acceptor electron will only experience the external magnetic field. Consequently, in the first half radical pairs the Larmor precession frequency of the donor electron, and thus the S-T mixing frequency, will be slightly higher than in the other half (due to the opposite contribution of the nuclear spin), hence the coherently formed triplet radical pairs will have their nuclear spins predominantly in the $|+\rangle$ state, leaving the singlet radical pairs with an equal and opposite nuclear polarization. While the

nuclear spins are thus sorted among singlet and triplet RPs, and unless there is a coherent mechanism like TSM producing net nuclear polarization, the expectation value $\langle I_z \rangle = \text{Tr}\{\rho I_z\}$ of the z-component of the single nuclear spin under consideration is still zero. Here ρ is the spin density matrix of the radical pair, describing the electron and nuclear spin degrees of freedom altogether. According to the traditional understanding of the radical-pair mechanism and the concomitant CIDNP signal generation, for this nuclear spin sorting to result in a diamagnetic ground state with net nuclear polarization, and in the absence of any coherent mechanism like TSM, at least one of the other two aforementioned incoherent mechanisms should be at play in order to observe nonzero CIDNP signals. Here we are going to show that this is not the case. To this end we will consider a magnetic Hamiltonian that *does not* exhibit the TSM mechanism, and we will also take $k_S = k_T$, i.e. we consider the singlet and triplet recombination rates of the RP to be equal, hence the differential decay (DD) mechanism is not operational.

To illustrate the above considerations, we consider a radical pair with one spin-1/2 nucleus *isotropically* coupled to the donor electron. In Fig. 2A we display the undisturbed spin state evolution brought about by the magnetic Hamiltonian $\mathcal{H} = A\mathbf{I} \cdot \mathbf{s}_1 + \omega(s_{1z} + s_{2z})$, where ω is the electron spin Larmor precession frequency and A the donor electron hyperfine coupling with the donor nuclear spin (in units of frequency). What is plotted in Fig. 2A is the spin state's singlet character, given by the expectation value of the singlet projection operator, $\langle Q_S \rangle = \text{Tr}\{\rho Q_S\}$, and assuming an RP with infinite lifetime. The nuclear spin, I_z , of singlet and triplet RPs, given by $\langle I_z \rangle^S = \langle Q_S I_z Q_S \rangle$ and $\langle I_z \rangle^T = \langle Q_T I_z Q_T \rangle$, is shown in Fig. 2B and 2C, respectively. The previous considerations regarding nuclear spin sorting are easily visualized by noting (dashed line L_1 in Figs. 2B and 2C) that although $\langle I_z \rangle^S \neq 0$ and $\langle I_z \rangle^T \neq 0$, the nuclear spin oscillations in the singlet and triplet subspace are exactly out of phase, so the net RP nuclear polarization $\langle I_z \rangle^S + \langle I_z \rangle^T = 0$. This is expected since the particular Hamiltonian $\mathcal{H}$ does not qualify for the TSM mechanism [22], while the differential decay mechanism (DD) involves the recombination reaction of RPs and so far is out of the discussion. Now, besides the coherent RP dynamics driven by $d\rho/dt = -i[\mathcal{H}, \rho]$ and producing the oscillations in Figs. 2A-2C, the spin-selective charge recombination of RPs has to be accounted for. This is done by a stochastic Liouville equation of the form $d\rho/dt = -i[\mathcal{H}, \rho] + \mathcal{L}(\rho)$, where $\mathcal{L}(\rho)$ is the reaction super-operator, written in terms of the singlet and triplet projection operators, Q_S and Q_T , and the singlet and triplet recombination rates, k_S and k_T , respectively. Recently, quantum measurement theory has been shown by Kominis [25, 26] to fundamentally describe the reaction dynamics of RPs, resulting in a new form of $\mathcal{L}(\rho)$, conceptually departing from the traditional theory used until now [27]. The main physical point of departure




FIG. 2: Time evolution of (A) $\langle Q_S \rangle$, (B) the nuclear spin of singlet RPs $\langle Q_S I_z Q_S \rangle$ and (C) the nuclear spin of triplet RPs $\langle Q_T I_z Q_T \rangle$, when the radical-pair spin state undergoes unceasing coherent S-T mixing driven by $d\rho/dt = -i[\mathcal{H}, \rho]$, where $\mathcal{H} = A\mathbf{I} \cdot \mathbf{s}_1 + \omega(s_{1z} + s_{2z})$. For this example it was $\omega = A/10$. The dashed line L_1 is a reminder of the fact that the singlet and triplet RP nuclear polarizations are exactly out of phase and sum up to zero. (D) Evolution of $\langle I_z \rangle$ according to the full density matrix equation of Kominis [26], with $k_S = k_T = 4A$. The absolute value of the nuclear polarization is shown in the right y-axis, while its enhancement relative to the thermal equilibrium polarization, on the order of 10^4 , is shown in the left y-axis. Qualitatively the same result (only higher by about a factor of 2) is obtained from the Jones-Hore theory [30]. In contrast, the traditional theory predicts $\langle I_z \rangle = 0$, exactly the same as the sum of (B) and (C) that applies to the case of no reaction. Note the different time scales in (A)-(C) and (D): since $k \gg \omega$, the build-up of $\langle I_z \rangle_{qc}$ in (D) takes place at the very beginning of the S-T mixing oscillations shown in (A)-(C).

is the fact that the very charge recombination process of RPs damps their coherent S-T mixing, i.e. it brings about an unavoidable intra-molecule decoherence in their spin state evolution. The traditional theory for the reaction operator of RPs, although successful in accounting for a large volume of experimental data, does not embody this decoherence process, neglecting which leads to several conceptual problems [28, 29]. In any case, in the example treated above, if the nuclear spin expectation value is calculated according to the full master equation of the traditional theory, taking $k_S = k_T$, it is found to be identically equal to zero, exactly as it is if only the Hamiltonian evolution is taken into account. According to the common wisdom, this should be perceived as an obvious result, since, with $k_S = k_T$ the differential decay mechanism (DD) cannot lead to any net nuclear polarization.

However, and this is the counter-intuitive new result presented here, if the new master equation derived by Kominis [26] is used, it is found that there actually is a non-zero RP nuclear spin polarization depicted in Fig. 2D. A similar result, different by just a factor of 2, follows from another variant of the quantum-measurement-based stochastic Liouville equation by Jones and Hore [30]. In this particular example, this nuclear polarization is about 10^4 times higher than the thermal equilibrium proton polarization (left y-axis of Fig. 2D) and roughly independent of ω as long as ω is relatively small (to be further discussed later). What is the source of this new kind of CIDNP signal? We will now go beyond the particular example considered above and provide the explanation for the general case of this new channel for photochemically enhancing nuclear polarizations: *the fundamental singlet-triplet decoherence underlying the radical-pair mechanism*.

To visualize this mechanism, suppose that at time t the single-nuclear-spin RP ensemble is in a state described by the density matrix ρ . Assuming a magnetic Hamiltonian *not* supporting any coherent generation of *net* RP nuclear polarization, as in the example considered previously, it will be $\langle I_z \rangle = \text{Tr}\{\rho I_z\} = 0$. This necessarily implies that $\langle I_z \rangle^S = -\langle I_z \rangle^T$. To parenthetically prove this, note that since I_z commutes with the electron spins $\mathbf{s}_1$ and $\mathbf{s}_2$, it also commutes with $Q_S = 1/4 - \mathbf{s}_1 \cdot \mathbf{s}_2$ and with $Q_T = 1 - Q_S$. Thus, since the two projection operators are orthogonal, i.e. $Q_S Q_T = Q_T Q_S = 0$, it will be $Q_S I_z Q_T = Q_T I_z Q_S = 0$. Furthermore, since $Q_S + Q_T = 1$, it will be $\text{Tr}\{\rho I_z\} = \text{Tr}\{\rho(Q_S + Q_T)I_z(Q_S + Q_T)\}$, and by using the previous relations it follows that indeed $0 = \langle I_z \rangle = \langle I_z \rangle^S + \langle I_z \rangle^T$.

Now, the main idea behind the new kind of CIDNP signal generation is this: the intramolecule S-T decoherence is equivalent to some RPs being *projected* to either the singlet or the triplet state, exactly due to the continuous quantum measurement of Q_S induced by the RP recombination dynamics, the measurement rate being $(k_S + k_T)/2$ [25]. The result of this measurement is either $Q_S = 1$, i.e. the RP is projected to the singlet manifold,

or $Q_S = 0$, i.e. the RP is projected to the triplet manifold. The probability for a singlet (triplet) projection is $\langle Q_S \rangle$ ($\langle Q_T \rangle$). Hence, at time $t+dt$ the state of the radical pairs will be (ignoring the RPs that recombine into neutral products during dt) a new mixture comprised of (a) radical pairs in the state $\rho + d\rho$, with $d\rho = -idt[\mathcal{H}, \rho]$, plus (b) the singlet-projected radical pairs described by $Q_S \rho Q_S$, plus (c) triplet-projected radical pairs described by $Q_T \rho Q_T$. The nuclear polarization of states (a) will still be zero, since again, the Hamiltonian $\mathcal{H}$ does not generate *net* nuclear polarization. The combined nuclear polarization of projected states (b) and (c), denoted by $\langle I_z \rangle^{\text{proj}}$, will be the *weighted* sum of singlet RP nuclear polarization, $\langle I_z \rangle^S$, and triplet RP nuclear polarization, $\langle I_z \rangle^T$, the sum being weighted by the respective projection probabilities, i.e.

$$\langle I_z \rangle^{\text{proj}} = \langle Q_S \rangle \langle I_z \rangle^S + \langle Q_T \rangle \langle I_z \rangle^T \quad (1)$$

Taking into account that $\langle I_z \rangle^T = -\langle I_z \rangle^S$ and using the completeness relation $\langle Q_T \rangle = 1 - \langle Q_S \rangle$, it follows that

$$\langle I_z \rangle^{\text{proj}} = \langle I_z \rangle^S (2\langle Q_S \rangle - 1) \quad (2)$$

Thus, if $\langle Q_S \rangle \neq \langle Q_T \rangle \neq 1/2$, more (when $\langle Q_S \rangle > 1/2$) or less (when $\langle Q_S \rangle < 1/2$) projections will take place to the singlet state than to the triplet, and $\langle I_z \rangle_t^{\text{proj}}$ will be non-zero, if of course $\langle I_z \rangle^S$ is non-zero.

We can now estimate the quantum measurement corrections to CIDNP, to be denoted by $\langle I_z \rangle_{\text{qc}}$, shown in Fig. 2D. Letting $k = k_S = k_T$, the rate of singlet or triplet projections (the "quantum measurement" rate) is just $(k_S + k_T)/2 = k$. We will focus on low magnetic fields and assume, as is the case most often, that $k > A, \omega$. Towards the estimate, consider the following three steps: (1) As the RPs start out in the singlet state, and the S-T mixing frequency $\Omega \ll k$, in the short reaction time of $1/k \ll 1/\Omega$ the singlet character will not have enough time to change appreciably, so it will be $\langle Q_S \rangle \approx 1$. (2) The electron spin of the donor molecule, which contains the one and only nuclear spin of the RP under consideration, will precess at frequency $|\omega \pm A|$ when the nuclear spin is in the $|\pm\rangle$ state. So at time t , the number of singlet RPs with the nuclear spin in the $|\pm\rangle$ state will be $N_{\pm} \approx \cos[(\omega \pm A)t]^2$. The nuclear polarization of singlet RPs at time $t = 1/k$ will thus be $\langle I_z \rangle^S = (N_+ - N_-)/(N_+ + N_-)$, easily seen to be

$$\langle I_z \rangle^S \approx -\frac{\omega A}{k^2} \quad (3)$$

And (3), S-T projections taking place at the rate k will be a source term for $d\langle I_z \rangle_{\text{qc}}/dt$. The source term, proportional to $\langle I_z \rangle^{\text{proj}}$, starts at $t = 0$ from zero and oscillates at a rate Ω while it is drained at a rate k , thus $d\langle I_z \rangle_{\text{qc}}/dt = k\langle I_z \rangle^{\text{proj}}e^{-kt} \sin \Omega t$, yielding at time $t \approx 1/k$ that $\langle I_z \rangle_{\text{qc}} \approx -\langle I_z \rangle^{\text{proj}}\Omega^2/k^2$. Putting everything together and using (2), we finally arrive at

$$\langle I_z \rangle_{\text{qc}} \approx \frac{\omega \Omega^2 A}{k^4} \quad (4)$$

For the particular example depicted in Fig. 2 it is (Fig. 2B) $\Omega \approx A/10$, hence according to the estimate (4) it is $\langle I_z \rangle_{\text{qc}} \approx 4 \times 10^{-6}$, in good agreement with the exact value shown in the right y-axis of Fig. 2D. Furthermore, the result (4) can be recast in terms of the thermal polarization $I_{\text{th}} = \hbar\omega/4\gamma k_B T$ (the expression for I_{th} is valid for $\hbar\omega \ll \gamma k_B T$), where $\omega = \gamma_e B$ is the electron Larmor frequency in the external magnetic field B , with $\gamma_e = 2\pi \times 2.8 \text{ MHz/G}$ being the electron gyromagnetic ratio and $\gamma = |\mu_e/\mu_p| = 658.5$ is the ratio of electron-to-proton magnetic moment. The enhancement factor $\langle I_z \rangle_{\text{qc}}/I_{\text{th}}$ is thus seen to be independent of ω and equal to

$$\frac{\langle I_z \rangle_{\text{qc}}}{I_{\text{th}}} \approx 10^3 \frac{(\Omega/0.01 \text{ ns}^{-1})^2 (A/0.1 \text{ ns}^{-1})}{(k/1 \text{ ns}^{-1})^4} \quad (5)$$

For ease of use, in the previous expression we provided the enhancement factor in terms of the S-T mixing frequency Ω (in units of 0.01 ns^{-1}), the hyperfine coupling A (in units of 0.1 ns^{-1}) and the recombination rate k (in units of 1 ns^{-1}). For typical hyperfine couplings of about 10 Gauss and recombination times on the order of 1 ns, it is seen that the quantum measurement corrections to chemically induced dynamic nuclear polarization amount to a significant enhancement of at least four orders of magnitude over the thermal polarization.

We note for the reader who would worry about angular momentum conservation that the RP's nuclear polarization is exactly balanced by an opposite and equal electronic polarization, i.e. the sum $\langle s_{1z} + s_{2z} + I_z \rangle = 0$ at all times, as it should be, since it is zero at time $t = 0$ (the initial state is $\rho_0 = Q_S/\text{Tr}\{Q_S\}$, which is a singlet state with zero nuclear polarization). The non-zero electronic polarization necessarily involves the admixture of the $|T_+\rangle$ and $|T_-\rangle$ triplet states, therefore this new CIDNP mechanism is a low-magnetic-field mechanism, because at high fields the $|T_{\pm}\rangle$ states are out of reach from the singlet state $|S\rangle$. How low a magnetic field depends on the recombination rate k , since the triplet RP decay broadens the triplet energy levels by about k , hence the mechanism is appreciable for magnetic fields B such that $B < k/\gamma_e$, i.e. when the Zeeman energy separation of $|T_{\pm}\rangle$ is within the broadening k . For recombination rates on the order of 1 ns^{-1} , the effect is appreciable for fields $B \leq 50 \text{ G}$.

In closing, the following comments should be made:

(1) Kaptein's rules provide a compact relation of the main features of the CIDNP signal to the system's parameters. For this case, the sign of the effect, i.e. whether it is emissive (+) or absorptive (-) is the same as the sign of the hyperfine coupling A .

(2) The new CIDNP effect reported here does not require any particular combination or fine-tuning of magnetic interactions, like the coherent TSM low-field effect analyzed in [24], but is rather general and shows up whenever there is a non-zero nuclear spin polarization of singlet RPs, $\langle I_z \rangle^S$, as in Fig. 2B. The rather

general requirement for this to happen is that the oscillation frequencies $\omega_m - \omega_n$ of the non-zero matrix elements $[Q_S]_{mn} = \langle m | Q_S | n \rangle$ must have terms linear in the hyperfine couplings. Here $|n\rangle$ and ω_n are the eigenvectors and eigenvalues of $\mathcal{H}$, with $n = 1, \dots, 4M$, where M is the nuclear spin multiplicity of the radical-pair.

(3) Experimentally, while NMR measurements at low fields suffer loss of sensitivity, recent methods of ultra-sensitive magnetometry using, SQUIDS [32, 33] or atomic magnetometers [34, 35], have demonstrated the ability to sensitively detect NMR signals at zero magnetic fields. So the detection of the new CIDNP signal analyzed here is rather straightforward.

(4) Although we considered the special case $k_S = k_T$, the new kind of CIDNP signal we predict will ubiquitously affect all sorts of radical pairs, including of course the ones with asymmetric recombination rates $k_S \neq k_T$, which predominantly occur in photosynthetic reaction centers [21]. We have chosen to deal with the special case $k_S = k_T$ and a magnetic Hamiltonian unable to support coherent generation of CIDNP signals in order to be able to focus exclusively on this fundamentally new CIDNP process. Establishing how this mechanism attributed to S-T decoherence affects, qualitatively and quantitatively, the cases where *there exists* a CIDNP signal according to the mechanism's traditional understanding (e.g TSM, DD or DR scenarios) is more involved an exercise that

will be undertaken elsewhere.

(5) We have considered the CIDNP signal in the RP state, and not in the diamagnetic ground state of the molecule in a closed reaction cycle as in Fig. 1B. However, it is easily seen that the same results hold for the ground state in the steady state under continuous excitation and a closed reaction cycle.

Concluding, this is the first step of a very promising research program, namely what are the particular repercussions of the recently unraveled fundamental quantum dynamics of the radical pair mechanism for understanding the dynamics of photosynthetic reaction centers. We have extended the fundamental understanding of CIDNP by pointing to a new and qualitatively different way to obtaining enhanced nuclear spin polarization. Interestingly, this polarization *naturally* turns out to be significant at low magnetic fields pertinent to natural photosynthesis. It stems from the random singlet or triplet projections taking place during the quantum evolution of radical pairs. S-T projections are the source of S-T decoherence fundamentally present in RP recombination reactions. Thus, an intriguing question raised is if and how the quantum mechanical process of spin decoherence affects the biological workings of photosynthetic reaction centers. In other words, what is the biological significance of earth-field CIDNP produced by S-T decoherence?

-
- [1] E. Collini et al., *Nature* **463**, 644 (2010).
 - [2] G. D. Scholes, G. R. Fleming, A. Olaya-Castro and R. van Grondelle, *Nature Chemistry* **3**, 763 (2011).
 - [3] P. Nalbach, D. Braun and M. Thwart, *Phys. Rev. E* **84**, 041926 (2011).
 - [4] M. M. Wilde, J. M. McCracken and A. Mizel, *Proc. R. Soc. A* **466**, 1347 (2010).
 - [5] A. Ishizaki and G. R. Fleming, *Proc. Natl. Acad. Sci. USA* **106**, 17255 (2009).
 - [6] G. Panitchayangkoon et al., *Proc. Natl. Acad. Sci. USA* **108**, 20909 (2011).
 - [7] M. Moshni, P. Rebentrost, S. Lloyd and A. Aspuru-Guzik, *J. Chem. Phys.* **129**, 174106 (2008).
 - [8] A. W. Chin, S. F. Huelga and M. B. Plenio, *Phil. Trans. Act. Roy. Soc.* **370**, 3638 (2012).
 - [9] M. Sarovar, A. Ishizaki, G. R. Fleming and K. B. Whaley, *Nature Physics* **6**, 462 (2010).
 - [10] S. G. Boxer, E. D. Chidsey and M. G. Roelofs, *Ann. Rev. Phys. Chem.* **34**, 389 (1983).
 - [11] Z. Schulten and K. Schulten, *J. Chem. Phys.* **66**, 4616 (1977).
 - [12] U. Steiner and T. Ulrich, *Chem. Rev.* **89**, 51 (1989).
 - [13] G. L. Closs and L. E. Closs, *J. Am. Chem. Soc.* **91**, 4549 (1969).
 - [14] R. Kaptein and J. L. Oosterhoff, *Chem. Phys. Lett.* **4**, 195 (1969).
 - [15] M. G. Zysmilich and A. McDermott, *J. Am. Chem. Soc.* **116**, 8362 (1994).
 - [16] T. Polenova and A. E. McDermott, *J. Phys. Chem. B* **103**, 535 (1999).
 - [17] S. Prakash et al., *J. Am. Chem. Soc.* **128**, 12794 (2006).
 - [18] A. Diller et al., *Proc. Natl. Acad. Sci. USA* **104**, 12767 (2007).
 - [19] E. Daviso et al., *J. Phys. Chem. C* **113**, 10269 (2009).
 - [20] E. Daviso et al., *Proc. Natl. Acad. Sci. USA* **106**, 22281 (2009).
 - [21] J. Matysik, A. Diller, E. Roy and A. Alia, *Photosynth. Res.* **102**, 427 (2009).
 - [22] G. Jeschke, *J. Chem. Phys.* **106**, 10072 (1997).
 - [23] G. Jeschke and J. Matysik, *Chem. Phys.* **294**, 239 (2003).
 - [24] G. Jeschke, B. C. Anger, B. E. Bode and J. Matysik, *J. Phys. Chem.* **115**, 9919 (2011).
 - [25] I. K. Kominis, *Phys. Rev. E* **80**, 056115 (2009).
 - [26] I. K. Kominis, *Phys. Rev. E* **83**, 056118 (2011).
 - [27] R. Haberkorn, *Mol. Phys.* **32**, 1491 (1976).
 - [28] I. K. Kominis, *Chem. Phys. Lett.* **542**, 143 (2012).
 - [29] I. K. Kominis, *Phys. Rev. E* **86**, 026111 (2012).
 - [30] J. A. Jones and P. J. Hore, *Chem. Phys. Lett.* **488**, 90 (2010).
 - [31] R. Kaptein, *Chem. Comm.* 732 (1971).
 - [32] R. McDermott et al., *Science* **295**, 2247 (2002).
 - [33] A. N. Matlachov et al., *J. Magn. Res.* **170**, 1 (2004).
 - [34] I. M. Savkov and M. V. Romalis, *Phys. Rev. Lett.* **94**, 123001 (2005).
 - [35] S. J. Xu et al., *Proc. Natl. Acad. Sci. USA* **103**, 12668 (2006).